

PATENT APPLICATION

MICROARRAYS FOR PERFORMING PROTEOMIC ANALYSES

Inventor: Deborah Charych
909 Taylor Street
Albany, CA 94706
U.S. Citizen

Eric Beausoleil
110 Gardenside Drive, #402
San Francisco, CA 94131
Citizen of Canada

Ronald N. Zuckermann
1116 Contra Costa Drive
El Cerrito, CA 94530
U.S. Citizen

Assignee: Chiron Corporation

MICROARRAYS FOR PERFORMING PROTEOMIC ANALYSES

CROSS-REFERENCE TO RELATED APPLICATIONS

5 This application claims priority from U.S. Provisional Application No. 60/209,711, entitled MICROARRAYS FOR PERFORMING PROTEOMIC ANALYSES, filed June 5, 2000, the disclosure of which is incorporated by reference herein for all purposes.

BACKGROUND OF THE INVENTION

10 The present invention relates to cell product analysis and materials. In one embodiment, the invention is directed to proteomic microarrays and methods of using them to conduct proteomic analyses.

15 In recent years, microarray technology has developed from a specialized sub-field into an important tool for basic and applied studies in molecular biology, microbiology, pharmaceuticals, agriculture, and many other biotechnologies. DNA microarray technology attempts to link the genome of an organism or cell to an expressed phenotype or protein function.

20 The overwhelming publication and patent literature on microarray technology describes arrays of DNA (or other forms of nucleic acid, such as cDNA or RNA), displayed on a solid surface such as a glass slide (often referred to as a "chip"). The arrayed DNA is typically in the form of short oligonucleotides (e.g., about 8 to 25 bases) or longer clones or PCR products (about 500 to 2000 bases). The former are typically synthesized on the solid support, whereas the latter are robotically "spotted" onto a solid support into an array format.

25 While there are reports of peptide and protein arrays on solid surfaces, these have received considerably less attention in comparison to DNA arrays. This is likely due to the inherent instability of these materials at interfaces, and in the presence of complex biological matrices. For example, it is well known, that many proteins denature upon contact with solid surfaces. Peptides, as well as proteins, are also subject to hydrolysis by any proteases that may be present in the biological sample being analyzed. In addition, peptide arrays are typically synthesized *in-situ* on solid surfaces using photolithographic methods. These techniques require the use of
30 expensive custom-made masks that must be designed and manufactured for each chip.

Furthermore, chemical characterization of surface-synthesized peptides is nearly impossible to perform due to the tiny amount of peptide generated.

Currently, the most common way of analyzing the proteome of biological samples employs two-dimensional ("2-D") gel electrophoresis. This method is problematic because the results are very sensitive to the experimental protocol (for example, development time of the gel as well as other parameters). Therefore, it is very difficult to get reproducible data from 2-D gels. Also, the sensitivity of the silver stain used in these gels is limited, and is less than that of the fluorescent labels used in microarray technologies.

Thus, there is an overwhelming need to develop effective microarray technology that is useful in a protein context. In many cases, functional pathways cannot be directly linked to a particular gene. Proteins often undergo a variety of post-translational modifications, interactions, or degradations that ultimately determine function. Even the seemingly simple evaluation of a protein's abundance cannot be directly correlated with the level of corresponding mRNA. The only solution is to evaluate the state of the cell, tissue or organism at the protein level. Therefore, a high throughput format that allows rapid display of protein differentials in complex mixtures such as cells, tissues, serum, etc., would provide a powerful counterpart and complement to DNA microarray technology.

SUMMARY OF THE INVENTION

To achieve the foregoing, the present invention provides peptidomimetic protein-binding arrays, their manufacture, use, and application. The protein-binding array elements of the invention include a peptidomimetic segment, an anchor segment and a linker segment connecting the peptidomimetic and anchor segment. The invention contemplates peptidomimetic array element library synthesis, distribution, and spotting of array elements onto solid planar substrates, labeling of complex protein mixtures, and the analysis of differential protein binding to the array. The invention also enables the enrichment or purification, and subsequent sequencing or structural analysis of proteins that are identified as differential by the array screen. Kits including proteomic microarrays in accordance with the present invention are also provided.

In one aspect, the invention pertains to an array of protein-binding agents stably attached to the surface of a solid support. The array includes a solid substrate having a substantially planar surface, and a plurality of different protein-binding agents bound to the substrate. Each of the protein-binding agents includes an anchoring segment stably bound to the substrate surface, a peptidomimetic protein-binding segment, and a linker segment connecting and separating the anchoring and peptidomimetic segments.

In another aspect, the invention pertains to a method of making an array comprising a plurality of different protein-binding agents stably associated with the surface of a solid support. The method involves preparing for bonding a solid substrate having a substantially planar surface, and contacting a plurality of different protein-binding agents with the substrate under conditions sufficient for the protein-binding agents to become bound to the substrate surface. Each of the protein-binding agents includes an anchoring segment stably bound to the substrate surface, a peptidomimetic protein-binding segment, and a linker segment connecting and separating the anchoring and peptidomimetic segments.

In another aspect, the invention pertains to a method of making an array comprising a plurality of different protein-binding agents stably associated with the surface of a solid support. The method involves generating a library of protein-binding agents in which each of the agents includes an anchoring segment stably bound to the substrate surface, a peptidomimetic protein-binding segment, and a linker segment connecting and separating the anchoring and peptidomimetic

segments. The protein-binding agents are distributed from the library into individual storage receptacles for each different protein-binding agent, a plurality of the different protein-binding agents are prepared for binding to a solid substrate, a solid substrate having a substantially planar surface for binding with a plurality of the different protein-binding agents is prepared, and an array is prepared as described herein.

A further aspect of the present invention pertains to a method of performing a differential binding assay. The method involves labeling proteins in a protein-containing biological sample solution, contacting an aliquot of the labeled protein-containing biological sample solution with an array as described herein, and analyzing the array to determine differential binding of proteins in the sample to protein-binding agents of the array.

Another aspect of the present invention pertains to a kit for use in performing a differential binding assay as described herein. The kit includes an array having a solid substrate having a substantially planar surface with a plurality of different protein-binding agents bound to the substrate. Each of the protein-binding agents includes an anchoring segment stably bound to the substrate surface, a peptidomimetic protein-binding segment, and a linker segment connecting and separating the anchoring and peptidomimetic segments.

A further aspect of the present invention pertains to a mixed array of protein-binding agents stably attached to the surface of a solid support. The array includes a solid substrate having a substantially planar surface, and a plurality of different protein-binding agents bound to the substrate. Each of the protein-binding agents includes an anchoring segment stably bound to the substrate surface, a peptidomimetic protein-binding segment, and a linker segment connecting and separating the anchoring and peptidomimetic segments. The array further includes a plurality of different antibodies bound to the substrate.

These and other features and advantages of the present invention will be presented in more detail in the following specification of the invention and the accompanying figures which illustrate by way of example the principles of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

Figs. 1A and 1B schematically depict the structure of a protein-binding agent array element and array portion, respectively, in accordance with one embodiment of the present invention.

5 Fig. 1C depicts the molecular structure a protein-binding agent linking segment composed of ethylene oxides in accordance with one embodiment of the present invention.

10 Figure 2 schematically depicts a process of making an array having a plurality of different protein-binding agents stably associated with the surface of a solid support in accordance with one embodiment of the present invention.

15 Figure 3 schematically depicts the molecular structures of six 12-mer peptoid array elements in accordance with one embodiment of the present invention.

20 Figures 4A and 4B schematically depict alternative modes of binding a protein-binding agent to a solid support in accordance with one embodiment of the present invention.

25 Figure 5 schematically depicts a process for conducting a differential protein binding assay in accordance with one embodiment of the present invention.

30 Figure 6 schematically depicts a various processes for identifying and characterizing proteins identified in accordance with embodiments of the present invention.

 Fig. 7A depicts the chemical formula for a NHS-LC-LC-biotin molecule used in a solution to coat aluminum slides to be used as a substrate in accordance with one embodiment of the present invention.

 Fig. 7B depicts a representation of an avidin-derviatized aluminum slide spotted with a biotinylated protein-binding agent in accordance with one embodiment of the present invention.

 Fig. 8 depicts a graph of results of size exclusion chromatography conducted to separate labeled protein from the unreacted dye prior to application of the labeled protein sample to a microarray in accordance with embodiments of the present invention.

Figs. 9A-9C depict scans of microarray chips bearing a library of 1,000 peptoid-based protein-binding agents used to prove the concept of the present invention.

5 Fig. 10 depicts biotinylated hexameric peptides (A through E) and the corresponding signal intensity obtained when the biotinylated peptides were spotted onto avidin-treated slides, blocked with casein, and probed with Cy5-labelled antiglu antibodies.

10 Fig. 11 depicts the dependence of signal strength on the mode of surface attachment of a peptide for two modes of surface attachment in accordance with the present invention.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212

DESCRIPTION OF SPECIFIC EMBODIMENTS OF THE INVENTION

The materials and associated techniques and apparatuses of the present invention will now be described with reference to several embodiments. Important properties and characteristics of the described embodiments are illustrated in the structures in the text and in the accompanying drawings. While the invention will be described in conjunction with these embodiments, it should be understood that the invention it is not intended to be limited to these embodiments. On the contrary, it is intended to cover alternatives, modifications, and equivalents as may be included within the spirit and scope of the invention. In the following description, numerous specific details are set forth in order to provide a thorough understanding of the present invention. The present invention may be practiced without some or all of these specific details. In other instances, well known process operations have not been described in detail in order not to unnecessarily obscure the present invention.

In this specification and the appended claims, the singular forms “a,” “an,” and “the” include plural reference unless the context clearly dictates otherwise. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention belongs.

Introduction

The present invention provides peptidomimetic protein-binding arrays, in addition to methods for their manufacture, use, and application. The protein-binding array elements of the invention include a peptidomimetic segment (an example of which is a peptoid), linked to a solid support using a stable anchor. In one embodiment, the invention provides peptidomimetic array element library synthesis, distribution, and spotting of array elements onto solid, planar substrates, labeling of complex protein mixtures, and the analysis of single protein binding or differential protein binding by contacting labeled pure protein or complex protein mixtures to the peptidomimetic array. Such analysis may lead to the identification of novel therapeutic targets involved in diseases such as cancer, HIV or diabetes. These novel targets can then be utilized in high throughput screening assays to find new drug candidates. The invention also enables the enrichment or purification, and subsequent sequencing or structural analysis, of proteins that are identified as differential by the array screen. The arrays can also be used to identify new synthetic ligands for proteins of interest (for protein purification or development of high throughput

assays) or synthetic antigens for antibodies of interest. They may also be used to find new enzyme inhibitors or other high affinity ligands for drug targets, or as initial scaffolds for new therapeutics. Kits including proteomic microarrays in accordance with the present invention are also provided.

5 1. Protein-Binding Agent Arrays

Peptidomimetic arrays in accordance with the present invention are composed of a number of different array elements comprising protein-binding agents attached to the surface of a solid support. The different protein-binding agent array elements each include an anchoring segment attached to the substrate surface, a peptidomimetic protein-binding segment, and a linker segment connecting and separating the
10 anchoring and peptidomimetic segments. A “peptidomimetic” as used herein refers to nonpeptide synthetic polymers or oligomers that detectably interact with proteins or receptors in a manner analogous to protein-protein or protein-peptide physical and/or chemical interactions under assay conditions. The anchoring segment is typically
15 attached to the substrate surface such that the association between the anchoring group and the substrate surface, and thus the concentration and density of the protein-binding agent on the substrate surface, is maintained during the processing to which the microarray is subjected under its normal operating conditions, for example, as described herein. The number of different chemical species of protein-binding agent
20 present on the surface of the array is at least 2, or at least 10, or at least 100, and can be much higher, generally being at least about 1,000, or at least about 5,000 to about 50,000, for example, between about 5,000 and about 10,000, as described further below.

Figs. 1A and 1B provide a representation of one such array element and an
25 array in accordance with the invention. In Fig. 1A, a protein-binding array element 100 includes three segments: a peptidomimetic segment 102, an anchor segment 104, and a linker segment 106. The array element is attached to a solid support or substrate 108 by a bond between its anchor segment 104 and the substrate surface 110. In some cases, the substrate may be composed of a plurality of layers 112a, 112b, 122c
30 forming a laminate. In Fig. 1B, a protein-binding array includes a plurality of different array elements 100, such as illustrated in Fig. 1A, attached to a planar substrate 108. Each of these components of the array is described in greater detail separately below.

A. Substrate

The solid support (Figs. 1A and 1B, element 108) employed in arrays in accordance with the present invention may vary greatly depending on the intended use of the product to be produced. The solid support may be any suitable material for binding the protein-binding agent that is also compatible with any analytical methods with which the array is to be used. Compatibility can be determined using methods and materials known to those having skill in the surface or materials chemistry arts. In one embodiment, the solid support comprises an impermeable, rigid material. Suitable materials include plastics, such as polymers, e.g. polyvinylchloride, polyethylene, polystyrenes, polyacrylates, polycarbonate and copolymers thereof, e.g., vinyl chloride/propylene polymer, vinyl chloride/vinyl acetate polymer, styrenic copolymers, and the like. Suitable materials also include glasses, such as those formed from quartz, or silicon; and metals (including alloys), e.g., gold, platinum, silver, copper, aluminum, titanium, chromium and the like.

As noted above, in many embodiments of interest, the support or substrate 108 will be a composite of two or more different layers of material, where the composition includes at least a base material, e.g., as represented by element 112a in Fig. 1A, and one or more surface coating materials, as represented by elements 112b and 112c in Fig. 1A. For example, one embodiment of interest includes a base material 112a, such as a glass, which is coated with a metallic layer 112b, e.g., gold or aluminum overcoated with silicon dioxide, or titanium overcoated with silicon dioxide, and a functionalized organic, e.g., amino-modified thiol or amino-modified silane, layer 112c. The planar solid support may be in the dimensions of a standard 3" x 1" microscope slide or in the shape of a 3" or 5" diameter circular wafer, for example. Other configurations will be apparent to those having skill in the surface or materials chemistry arts.

The solid support material or substrate may have a variety of different configurations, depending on the intended use of the material. Thus, in the broadest sense the support or base material may be in the form of a plate, sheet, cone, tube, well, bead, nanoparticle (e.g., about 5-100 nm), wafer, etc. In some embodiments, the base support material is one that has at least one substantially planar surface, e.g., as found on a plate, slide, sheet, disc, etc. In these embodiments, supports having an overall slide or plate configuration, such as a rectangular or disc configuration are employed.

In the planar, rectangular embodiments of the above-described slides, the length of the support will generally be at least about 1 cm and may be as great as about 40 cm or more, but will usually not exceed about 30 cm and may often not exceed about 20 cm. The width of support will generally be at least about 1 cm and may be as great as about 40 cm, but will usually not exceed about 30 cm and will often not exceed about 20 cm. The height of the support will generally range from about 0.01 mm to about 10 mm, depending at least in part on the material from which the rigid substrate is fabricated and the thickness of the material required to provide the requisite rigidity. Of particular interest in many embodiments are supports having the dimensions of a standard microscope slide. One typical substrate size is about 2.54 cm x 7.62 cm and about 1-2 mm thick. However, any suitable dimensions can be employed.

Where the support is a bead, nanoparticle or microparticle, the diameter of the support typically ranges from about 5 nm to about 1000 μ , particularly from about 10nm to about 500 μ .

The protein-binding agents can either be attached directly to the inorganic solid surface of layers 112a or 112b (as illustrated and described in more detail below with reference to Fig. 4A) or attached using the functionalized organic layer of 112c (as illustrated and described in more detail below with reference to Fig. 4B). In the latter case, the non-substrate-bound termini of the organic molecules in the layer are functionalized with a reactive group that can attach to the anchoring group of a subsequently bound protein-binding agent. Suitable terminal reactive groups may be, for example, maleimide, hydrazide, aminooxy, an activated ester such as N-hydroxysuccinimide, anhydride, aldehyde, disulfide, thiol, azide, phosphine or avidin, streptavidin, neutravidin or other altered forms of the protein avidin that bind biotin, depending upon the anchor's functional group.

In those embodiments in which the surface of the base support material, such as glass, is coated with a thin layer of a metal, such as aluminum, gold, or titanium, the thickness of the metal layer will generally range from about 300 Å to about 10,000 Å, more particularly from about 750 Å to about 2,000 Å, and still more particularly from about 1,000 Å to about 1,500 Å. The metal layer may be deposited on the substrate surface using any suitable protocol, including e-beam deposition, vapor deposition, sputtering and the like, as are known to those of skill in the art. An adhesion metal layer may be present between the metal layer and the substrate, where adhesion metals of interest include titanium, chromium, and the like. When present,

the adhesion metal layer will typically range in thickness between about 5 Å and about 100 Å, usually between about 25 Å and about 75 Å and in many embodiments will be about 50 Å. In some embodiments, the above-described adhesion layer can be a molecular adhesion layer. Examples of materials suitable for forming molecular adhesion layers in accordance with the present invention include mercaptopropyltriethoxysilane, and other mercaptoalkoxysilanes, such as mercaptopropyltrimethoxysilane, mercaptopropyltrichlorosilane, or other chain lengths such as mercaptohexyltriethoxysilane and other mercaptohexylalkoxysilanes, as are known in the art. Where the adhesion layer is a molecular adhesion layer, the thickness of the adhesion layer typically ranges from about 5 Å to about 50 Å.

The anchoring group of a protein-binding agent may be used to directly bond the protein-binding agent to an inorganic, e.g., metal or glass, substrate surface. For example, a thiol anchoring group may be used to bond directly to metals such as gold silver, copper or platinum without an intervening functionalized layer (e.g., maleimide/amine/thiol). Or, if the anchor group is an activated silane (i.e., modified to be reactive with other organic species, for example, hydrolyzable silane or modified silane that can condense with hydroxyls or other silanes to form siloxane bonds), e.g., chlorosilane or an alkoxysilane, the protein-binding agent can be directly bonded to glass or other oxide surfaces, such as titanium oxide, silicon oxide or aluminum oxide. Other activated silanes for this purpose include triethoxysilane, trichlorosilane, trimethoxysilane or other trialkoxysilanes. Within each of the classes, chain lengths may be from three atoms up to about eighteen atoms, for example.

As will be described in more detail below, in the case of bonding of the protein-binding agent directly to a substrate's inorganic surface, the protein-binding agent's linking segment (between the peptidomimetic and the anchor) should be long enough to keep the peptidomimetic sufficiently far from the hard substrate surface that the surface does not interfere with protein binding occurring (subsequently) at the peptidomimetic. The length can be from about 2 atoms to about 200 atoms, or about 2 Angstroms to about 300 Angstroms, for example. A typical linking segment may have a backbone of between about 2 to about 30 atoms, preferably about 6 to about 12 atoms. The linking segment may be composed of, for example, suitably derivatized aliphatic chains (e.g., aminoalkanoic acids, such as aminohexanoic acid), ethylene oxides (e.g., such as shown in Fig. 1C), sulfoxides, or "non-binding" (orthogonal) short peptoid or peptide elements that remain constant for each element of the array (e.g., a glycine heptamer), or some combination of these components.

As noted above, the protein-binding agents may also be attached using a functionalized organic layer (Fig. 1A, element 112c), such as an amino-modified thiol (aminothiol) (for use with some metal substrate surfaces) or an amino-modified silane (aminosilane) (for use with glass, metallic or other oxide substrate surfaces). Where such an organic layer is used, the termini of the organic molecules of the layer are functionalized with a reactive group that can stably attach to the substrate surface at one end, and to the anchor segment of a subsequently-bound protein-binding agent in accordance with the present invention at another end. Suitable terminal groups include, for example, maleimide which can form a covalent bond upon reaction with a thiol presented in the anchor. A further advantage of such synthetic anchoring systems is their stability, conferring long shelf life. Other possible terminal groups on the substrate include hydrazide, which can react to form covalent bonds with aldehyde or ketone moieties in the anchor; aminooxy, which can react to form covalent bonds with ketone moieties in the anchor, aldehyde, disulfide, thiol, azide, phosphine.

In another implementation, proteins such as avidin, streptavidin, or other analogs may be used as a coating for the solid support. In this case, biotin is used as the anchoring moiety of the protein-binding agent to form a stable, non-covalent bonding complex with avidin on the surface. Without wishing to be bound by any particular theory of action, it is believed that this format displays the protein-binding agent in a particularly biocompatible environment with desirable distances between display molecules and their neighbors, and desirable distances between display molecules and the solid surface. Other protein-coatings can also be employed, provided they sufficiently bind a small molecule anchoring group. These may include anti-digoxigenin / digoxigenin, anti-dinitrophenol / dinitrophenol, or many other protein / small molecule pairs known in the art. Avidin / biotin is of particular value because of its extremely stable binding interaction. In addition, substantial signal increases have been observed for biotin / avidin immobilization compared to other suitable immobilization techniques in accordance with the present invention, for example 1000X higher than thiol / maleimide. Suitable synthetic macromolecules may also be used as mimics of such protein spacers. These may include high molecular weight polymers such as polyethyleneimines, dendrimers, polyacrylic acids, polylysines and the like.

Of course, other suitable binding combinations of this character (namely, as noted above, sufficiently stable to maintain the bond during the processing to which the microarray is subjected under its normal operating conditions) are also possible. In addition, those having skill in the surface chemistry arts will understand that the

above-listed anchoring groups on the peptoid can function as reactive groups on the surface, and the reactive surface groups can also function as a peptidomimetic anchoring groups.

In one embodiment of the present invention, aluminum slides may be coated with a layer of silicon dioxide or silicon monoxide having a thickness of between about 500 Å to about 2,000 Å. The thickness is chosen to roughly correspond to 1/4 the wavelength of the emission or excitation light. A layer of an aminoalkyl trialkoxysilane, such as aminopropyl triethoxysilane (APS), trichlorosilane, trimethoxysilane, and any other trialkoxysilane is coated on the surface of the oxide. In addition, other amino-silanes could also be used, for example, compounds having longer alkyl groups, such as octyl, decyl, hexadecyl, etc., that may form more ordered silane layers as will be appreciated by those having skill in the surface chemistry arts. The thickness of this silane layer may be from about 3 Å to about 100 Å, more preferably about 5 Å to about 50 Å, even more preferably about 7 Å to about 20 Å. One suitable example is an APS layer that is about 7 Å thick. The amino-modified Al surfaces may be functionalized with a reactive group that will bind to the anchor functional group on a protein-binding agent. In one embodiment, the functional group may be maleimide. In another embodiment, the functional group may be a whole protein such as avidin. An implementation of an avidin-presenting substrate is described with reference to Figs. 7A and 7B in Example 3, below.

In another embodiment, a gold-surfaced substrate slide (or a slide surfaced with any metal capable of forming metal-thiol bonds, such as Ag, Pt, or Cu) may be coated with an aminothiols layer that is functionalized with a group that will bind to the anchor functional group on a protein-binding agent, such as maleimide. However, the anchor functional group and the surface-bound reactive group should be chosen to have orthogonal reactivities. As noted above, these anchor and substrate-bound groups can be interchanged as will be readily understood by those skilled in the art.

In general, the functionalized organic layer 112c is characterized by having a substantially uniform hydrophilic surface. By uniform is meant that the surface of the layer includes substantially no irregularities, such as gaps, pinholes, etc. The thickness of the layer 112c may vary considerably, where the thickness may be less than about 5,000 Å, usually less than about 2,000 Å, but can be much lower, particularly from about 5 Å to about 100 Å, or between about 20 Å to about 50 Å.

In at least those embodiments where the support material includes a metallic layer beneath a functionalized organic layer, e.g., where the support material is an

amino-modified thiol layer on a gold coated microscope slide as described above, the thickness of the organic layer is chosen such that the layer 112c is at least sufficiently thick to separate any fluorescently labeled moiety that may be present to the surface a sufficient distance from the metallic layer on the substrate surface such that significant signal quenching (i.e., signal quenching of sufficient magnitude to effectively preclude meaningful detection of the signal) does not occur. In these embodiments, the functionalized organic layer may have a thickness that is at least about 30 Å, usually at least about 50 Å and more usually at least about 100 Å. Alternatively, in cases where quenching does occur, the slide may be treated with a high salt solution such as 0.75 M sodium chloride, 0.085 M sodium citrate (*see. e.g.*, PCT Publication No. WO 01/01142).

In those embodiments where the array is employed with fluorescently labeled target, as described in greater detail below, the thickness of the functionalized amino-modified thiol layer may be chosen to provide for maximum amplification of the emitted and reflected signals. See e.g., U.S. Patent No. 5,055,265, the disclosure of which is herein incorporated by reference. In such embodiments, the thickness of the organic layer will be about ¼ of the wavelength of the emitted light from the label. While the exact thickness of the organic layer will vary depending on the particular label with which it is to be employed, the thickness generally ranges from about 50 Å to about 300 Å, usually from about 100 Å to about 200 Å and more usually from about 125 Å to about 150Å.

In one embodiment, the functionalized amino-modified thiol layer 112c includes at least one self-assembled monolayer (SAM). As such, the functionalized organic layer may include one or more different self-assembled monolayers grafted sequentially onto each other, where when the layer includes more than one self-assembled monolayer (*see. e.g.*, PCT Publication No. WO 01/01142).

B. Protein-Binding Agents

As noted above, the protein-binding agents of the present invention are composed of three segments: a peptidomimetic, an anchor, and a linker connecting the peptidomimetic and the anchor.

A “peptidomimetic” as used herein refers to nonpeptide synthetic polymers or oligomers that detectably interact with proteins or receptors in a manner analogous to protein-protein or protein-peptide physical chemical interactions under assay conditions. Peptidomimetics are generally protease-resistant, and include, for

example, oligomeric species such as peptoids, beta-peptides, and others as described in A.E. Barron & R.N. Zuckermann, *Bioinspired polymeric materials: in-between proteins and plastics*, Curr Opin Chem Biol 1999, 3(6):681-7 and K Kirchenbaum, R.N. Zuckermann and K.A. Dill, *Designing polymers the mimic biomolecules*, Curr Opin Chem Biol 1999 9(4):530-5, each of which is incorporated by reference herein in its entirety for all purposes, and constrained cyclic molecules such as cyclic peptides and heterocycles. In some cases, the peptidomimetic may also mimic the folding of natural proteins. Peptidomimetics in accordance with the present invention comprise generally no more than about 500 mers, more particularly no more than about 100 mers, typically no more than about 50 mers, and usually about 10 to about 20 mers, more particularly about 12 to about 16 mers. Given the parameters provided herein, one of skill in the art would be able to choose an appropriate peptidomimetic length for a given application without undue experimentation.

Examples of peptidomimetic libraries and their design and synthesis may be found in Fahad Al-Obeidi, et al., *Peptide and Peptidomimetic Libraries*, Molecular Biotechnology 1998, 9: 205-223; Victor J. Hruby et al., *Synthesis of oligopeptide and peptidomimetic libraries*, Curr Opin Chem Biol 1997, 1:114-119; and Amy S. Ripka et al., *Peptidomimetic Design*, Curr Opin Chem Biol 1998, 2:441-452, incorporated by reference herein in its entirety for all purposes.

In a preferred embodiment of the present invention, the peptidomimetic segment of the protein-binding agent is a peptoid. The term "peptoid" as used herein refers to polymers comprising N^α-substituted amides as described in U.S. Patents Nos. 5,831,005, 5,877,278, 5,977,301, 5,871,387, 5,986,695, and 5,719,049; and in co-pending U.S. Patent Applications Serial Nos. 08/340,073, 08/836,167, 08/920,205, 08/126,539, 08/277,228, 08/454,511, 08/485,106, 09/132,828, 08/484,923, 09/704,422; and PCT Publications Serial Nos. 96/15143 and 93/09117, each of which is incorporated herein by reference in its entirety and for all purposes.

The anchoring segment provides for the stable attachment of the array element to the solid surface. This functional group is chosen to be reactive towards a complementary group that is displayed on the solid surface and orthogonal to (*i.e.*, substantially non-reactive with) sidechains present in the ligand. An important feature of the anchoring group is that its reaction to the surface be sufficiently facile so that it is complete within the average lifetime of a droplet that is deposited by the robotic array spotter onto the surface. For example, if the surface displays a maleimide, a suitable anchoring group is a thiol and approximately 15-20 minutes at about 60%

humidity are required for completion of the binding reaction before the approximately 10 nL drop evaporates. Of course, droplet lifetime varies with temperature, humidity and other conditions, allowing more or less time for the reaction to take place. As noted above, other surface display/anchor combinations are possible, including a hydrazide surface group with an aldehyde or ketone anchoring group. Or, the anchoring group can also be a biotin molecule, that will attach strongly (yet non-covalently) to a surface that displays an avidin protein.

sub B1
The anchoring group may be attached to the peptidomimetic at either end (in the case of a peptoid, at either the C- or N-terminus). It can be attached either as a submonomer (e.g., a substituted amine), or as a modification of a peptidomimetic (e.g., peptoid) side chain after synthesis. Alternatively, in another example, the anchoring group may be present as a linker connecting the peptoid to a beaded solid support upon which it is synthesized. This linker may be cleaved from the resin along with the peptoid, to provide a readily available anchoring group. Some examples are described in co-pending U.S. Patent Application Serial No. _____ (Attorney Docket No. 16708.001), titled *Peptoids Incorporating Chemoselective Functionalities*, filed April 6, 2001, which is incorporated herein by reference in its entirety and for all purposes.

The linking segment of the protein-binding agent molecule is chosen to provide for separation between the solid surface and the peptidomimetic segment sufficient to facilitate interaction between the peptidomimetic and the components of the analyte solution (solution with which the microarray will be contacted), for example by providing separation between the substrate surface and the peptidomimetic so that the surface does not interfere with protein binding occurring (subsequently) at the peptidomimetic access for a protein binding site to the peptidomimetic ligand displayed on the surface, especially for proteins with deep binding pockets. The linker may also serve to separate the peptidomimetics on the surface from each other, thereby mitigating possible steric hindrance between the ligand and protein binding pocket. A typical linking segment may have a backbone of between about 2 to about 200 atoms, preferably about 6 to about 30 atoms. The linking segment may be composed of, for example, aliphatic chains (e.g., aminoalkanoic acids, such as aminohexanoic acid), ethylene oxides, sulfoxides, or "non-binding" ("orthogonal") short peptoid or peptide elements that remain constant for each element of the array, or some combination of these components. In one embodiment, a 2-carbon linker may be used. In another embodiment, three ethylene oxides may be used. An example of a non-binding short peptoid is a 2-mer to 12-mer,

for example a 4-mer of methoxyethyl side chains that remain constant for each protein binding agent, while the peptidomimetic segment is variable. A suitable orthogonal peptide linker is a 2-mer to 12-mer, for example a 5-mer of glycine.

In some embodiments, where an organic layer is present on the substrate surface, a spacer functionality may be built into the organic layer. In such cases, the spacer in the substrate surface layer and the linker in the protein-binding agent may collectively contribute to the desired spacing of the peptidomimetic from the hard substrate surface. Further, the mitigation of steric hindrance may also be achieved by the selection of a surface coating that presents a particularly biocompatible format, such as an avidin protein (coupled with a biotin anchoring group on the protein-binding agent).

Fig. 3 illustrates an example of a small set of array elements 300 wherein the peptidomimetic protein-binding segments 302 are 12-mer peptoids, the linker 304 is a short aliphatic chain, and the anchoring group 306 is a thiol. The peptidomimetic segments depicted in Fig. 3 illustrate a subset of the range of affinity properties achievable using peptoids as the peptidomimetic segment, in accordance with one embodiment of the present invention

As mentioned above, the number of different types of binding agents present on the surface of the array is at least two. By "different", it is meant that the sequence of monomeric units between two different peptidomimetics of two different protein-binding agents is not the same. While the number of different species of protein-binding agents present on the surface of the array is at least 2, at least about 10, at least about 50, or at least about 100, it is typically much higher, generally being at least about 1,000 usually at least about 5,000 and more usually at least about 10,000. The number may be as high as 500,000 or higher, but typically does not exceed about 100,000 and usually does not exceed about 50,000.

The surface regions surrounding the protein-binding array elements may be modified so as to minimize background non-specific binding of proteins, allowing complex samples (e.g., lysates or serum) to be examined in a single step. The surface may be blocked chemically with hydrophilic termini such as alcohols, carbohydrates, amino acids, sulfoxides, acrylamides, or ethers. Examples of alcohol terminal groups are, for example, mercaptoethanol, mercaptohexanol, mercapto-octanol, in the case of a maleimide-treated surface. These block the unreacted maleimides on the surface. The surface can also be blocked using proteins such as solutions of 1% - 10% bovine serum albumin ("BSA") or human serum albumin ("HSA") in phosphate buffered

saline, or 1% - 10% non-fat dry milk or 1% - 10% casein, gelatin or other suitable blocking protein. In some cases, the addition of detergents to the blocking protein solution is advantageous. For example, the addition of 0.01% - 0.5% Tween-20 or Triton X-100 (particularly 0.05%), 0.1 - 2% SDS as are well known in the assay development or surface chemistry arts.

C. General Features of the Array

Typically, the array is characterized by having a plurality of protein-binding agent spots on a solid substrate, where each spot is characterized by having one or more, usually a plurality, of identical binding agents bound to the support surface.

The number of distinct spots on the surface of the array may or may not be the same as the number of different protein-binding agents on the array, e.g., the same protein-binding agent may be presented in two or more spots on the array surface. In one embodiment, each protein-binding agent is presented in duplicate in the array. Depending on the nature of the binding agents, the size of the support surface, the methods of fabrication and the intended use of the array, the number of distinct spots on the array surface may vary greatly. Where the support surface has the dimensions of a standard microscope slide (about 3" x 1"), the number of spots on the support surface will typically be at least about 3,000, usually at least about 6,000 and more usually at least about 10,000 - 50,000. The number may be as high as 100,000 or higher, but typically does not exceed about 75,000 and usually does not exceed about 50,000.

The diameter of each spot will typically range from about 100 μm to about 300 μm , usually from about 200 μm to about 300 μm . The space between any two given spots will generally be between about 1 μm and about 50 μm . The density of the spots generally ranges from about 1 to about 5,000 spots/ cm^2 , usually from about 100 to 2,000 spots/ cm^2 . Typically, the spots are arranged across the surface of the spacer layer in the form of a pattern. The pattern may be in the form of organized rows and columns of spots, e.g., a grid of spots, across the substrate surface, a series of curvilinear rows across the substrate surface, e.g., a series of concentric circles or semi-circles of spots, and the like. To further increase density, the spots may also be hexagonally arranged. Still other arrangements of spots are within the scope of the present invention.

2. Methods of Making the Protein-Binding Agent Arrays of the Subject Invention

The arrays of the subject invention may be prepared using any convenient protocol. One protocol of interest involves 1) the procurement of a solid support having a surface activated for binding of a protein-binding agent; and 2) contact of two or more different protein-binding agents with the support surface under conditions such that the protein-binding agents become stably associated with the support surface.

A. Solid Support Fabrication

The solid support may be fabricated using any convenient methodology, which methodology will vary depending the particular nature of the solid support. In accordance with one embodiment of the invention a glass support is coated with a layer of metal, e.g., aluminum or gold. To prepare a solid support of glass coated with a metal layer, the surface of the glass is coated with a thin layer of the metal, e.g., gold, silver, platinum, copper, titanium, or aluminum, etc. in a thickness as described above. The metal layer may be deposited on the substrate surface using any convenient protocol, where suitable protocols include e-beam deposition, vapor deposition, sputtering, and the like, and are known to those of skill in the art. See e.g., Moteshari et al., J. Am. Chem. Soc. (1998) 120:1328-1336; Bain et al., J. Am. Chem. Soc. (1989) 111:7155-7164; Lee et al. Langmuir (1998) 14:6419-6423; Folkers et al., Langmuir (1992) 8:1330-1341. Where convenient, an adhesion metal layer may be present between the metal layer and the substrate, where adhesion metals of interest include titanium, chromium, and the like, deposited in a thickness as described above. In addition, oxide overlayers such as silicon dioxide or silicon monoxide may be deposited by e-beam or sputtering deposition on top of the metallic layer.

In one example, following preparation of a gold substrate, if the protein-binding agent's anchoring group is a thiol and the linking group is sufficiently long (as explained above), arrays in accordance with the present invention can be formed by spotting thiol-displaying protein-binding agents onto bare, clean gold. The gold surface of the substrate may be cleaned using a chromic acid cleaning solution (e.g., chromium oxide or sodium dichromate in sulfuric acid, for example Nochromix, available from Fisher (50-80% Sodium dichromate in 12 N sulfuric acid)) and rinsed with HPLC-grade water. This has the advantage of reducing the number of surface functionalization steps.

In another embodiment, where the protein-binding agent's anchoring group is a functionalized modified silane (e.g., mono-, di- or tri- functional silanes, such as chlorosilane, an alkoxysilane, dichloro or dialkoxo silane, or trichloro or trialkoxy silane), the anchoring group can be attached directly to oxide-containing surfaces such as glass, titanium oxide or aluminum oxide. Arrays in accordance with these
5 embodiments of the present invention can be formed by spotting active silane-
displaying protein-binding agents onto the oxidized substrate surface. A glass
substrate has surface hydroxyls available for this binding. Metal substrate surfaces
may be oxidized, for instance by thermal or chemical treatment. For example,
10 aluminum may be oxidized electrochemically, thermally or chemically (e.g., with
 H_2O_2), as is well known in the art. In addition, a silicon dioxide or titanium oxide
layer can be deposited on the aluminum. An oxide may also be present as a native
thin layer, such as occurs with aluminum or silicon. This native oxide may then be
derivatized by the amino-silane.

15 In still another embodiment of the invention, a metal, e.g., gold or aluminum,
substrate surface is first functionalized with functionalized organic molecules that
form ordered monolayers. The termini of the organic molecules are functionalized
with a reactive group that can attach to a suitable anchoring group of a protein-
binding agent. Suitable terminal groups may be, for example, maleimide, hydrazide,
20 aminooxy, an activated ester such as N-hydroxysuccinimide, anhydride, aldehyde,
disulfide, thiol, azide, phosphine or avidin, depending upon the anchor's functional
group.

25 In one embodiment the functionalized organic molecules are amino-modified
thiol molecules. In order to functionalize the metal surface with the thiol molecules,
the substrate having the metal surface may be dipped in a solution of the amino-
modified thiol; the amino-modified thiol solution then may be deposited onto the
surface of the substrate. Other convenient protocols may be employed. Typically, the
gold substrate is immersed into the amino-modified thiol solution under conditions
and for a sufficient period for the amino-modified thiol molecules to assemble into a
30 monolayer on the substrate surface. The temperature at which contact is carried out
typically ranges from about 10°C to about 100°C, usually from about 15°C to about
80°C. Contact is maintained for a sufficient period of time for the self-assembled
monolayer to form on the gold surface, where contact is typically maintained for at
least about 20 minutes, usually at least about 4 hours, and more usually at least about
35 16 hours.

Alternatively, the amino thiol can also be deposited by vapor deposition in a vacuum oven or by spin coating. For example, a 1 - 10% (e.g., 5%) solution of thiol in a volatile solvent such as isopropanol, methanol, THF may be prepared. The slides may be spun at about 1000 – 8000 rpm (e.g., 5000 rpm) to provide an even deposition of the thiol. Then, a heterobifunctional molecule (e.g., succinimidyl 4-(N-maleimidomethyl)-cyclohexane-1-carboxylate (SMCC) or LC (long-chain) -- SMCC) that is an activated ester on one end and a maleimide on the other is contacted with the amino group to create the maleimide-terminated surface. Other heterobifunctional cross-linkers could also be used with different length spacers (e.g., a long ethylene oxide spacer (e.g., 2 – 4 units) between an NHS ester on one end, and a maleimide on the other end. As noted above, the spacer on the substrate serve the same purpose as and the “linker” in the protein-binding agent.

In another embodiment, Aluminum slides may be used. Aluminum metal may be deposited by e-beam deposition onto a clean glass substrate. The aluminum is then overcoated by silicon dioxide (SiO_2) or silicon monoxide in a thickness that is the same or thinner than $1/4$ the wavelength of the emission or excitation light. Oxide thicknesses of about 600 to 1000 Angstroms may be used as these eliminate the need for thinning the oxide prior to performing binding experiments. The aluminum / oxide surface may be treated with a amino-modified silane. For example, aluminum slides freshly coated with a 800 – 1,400 Angstrom layer of silicon dioxide, may be dipped immediately into a bath of 3% - 40% aminopropyl triethoxysilane in isopropanol, that has been previously filtered through an 0.2 μM filter membrane, and silanized for up to 1 hour, followed by rinsing and drying. Alternatively, aluminum slides coated with silane (APS) are available from Amersham-Pharmacia, Amersham, England (and described in International Patent Application No. WO 98/53304). In some cases, such commercially available slides may require thinning of the oxide layer to between about 200 Å and about 1,000 Å, more particularly between about 900 Å and about 1,000 Å, prior to performing a binding experiment to improve signal-to-noise ratios. The final amino-modified Al surfaces may be functionalized with SMCC to render a surface that presents maleimide functional groups. Here again, the silane may also be vapor deposited or spin coated, as described above. For example, a 1 - 10% (e.g., 5%) solution of silane in a volatile solvent such as isopropanol, methanol, THF may be prepared. The slides may be spun at 1,000 – 8,000 rpm (e.g., 5,000 rpm) to provide an even deposition of the silane. Then, a heterobifunctional molecule (e.g., succinimidyl 4-(N-maleimidomethyl)-cyclohexane-1-carboxylate (SMCC or LC-SMCC)) that is an activated ester on one end and a maleimide on the other is contacted with the amino group to create the maleimide-

terminated surface. Other heterobifunctional cross-linkers could also be used with different length spacers (e.g., a long ethylene oxide spacer (e.g., 2 – 4 units) between an NHS ester on one end, and a biotin or maleimide on the other end. As noted above, the spacer on the substrate serve the same purpose as and the “linker” in the protein-binding agent.

Further in accordance with this embodiment, it has been found that amplification of the fluorescent signal used in assays conducted with microarrays in accordance with the present invention may be enhanced by etching the commercially available functionalized and spotted aluminum slides (Amersham) to reduce the oxide layer thickness to about 200 Å to about 900 Å, preferably about 500 Å. For example, an etch solution of about 0.1 – 0.2 % SDS / 5X SSC (0.75 M NaCl/0.085 M sodium citrate) and optionally 5mM EDTA may be applied at about 50 – 80 degrees, preferably about 60 °C for about two to four hours. As noted above, manual deposition of a silicon dioxide layer of a suitable thickness may eliminate the need for the thinning of the slide.

As noted above, where the substrate surface has an organic self-assembled monolayer, following self assembly of the initial monolayer, one or more additional monolayers may be grafted onto the initial monolayer, where the additional layer(s) are made up of molecules, e.g. alkyls, having functionalities that provide for their covalent attachment to the surface functionalities of the initially deposited monolayer, e.g. amino functionalities where the initially deposited monolayer is characterized by the presence of carboxy functionalities. Alternatively, multi-component spacers can be constructed prior to attachment to the surface.

B. Synthesis of the Protein-Binding Agents

As noted above, protein-binding agents in accordance with the present invention are composed of three segments: a peptidomimetic, an anchor, and a linker connecting the peptidomimetic and the anchor.

Peptidomimetics may have a variety of nonpeptide polymeric structures and characteristics as long as they can mimic the biological effects of natural peptides. References to peptidomimetic structures have been provided above. In one embodiment of the present invention, the peptidomimetic segment of the protein-binding agent is a peptoid.

Libraries of peptoids may be synthesized using robotic solid-phase synthesis techniques, such as those developed by Chiron Corporation of Emeryville, CA. The diversity of the library can be controlled by the nature and arrangement of the amine submonomers (bromoacetic acid and substituted amines) used in the peptoid synthesis, as described in above-incorporated U.S. patent documents. As illustrated in Fig. 2, libraries may be prepared according to a process 200 using mix and split synthesis 202, or parallel synthesis, such that there is a multiplicity of peptoids synthesized, but only one species of peptoid on a given bead (i.e., each bead has one unique compound, repeated many times). The amount of compound per bead may range from about 1 – 100 nanomoles, with 20 – 40 nanomoles being most typical. About 40 nanomoles per bead are commonly obtained. This process is further described in U.S. patent Application No. Serial 09/306,700, the disclosure of which is incorporated by reference herein in its entirety and for all purposes. The synthesized peptoids, still bound to resin, may then be distributed to 96-well plates (or other multi-well plates, such as 384-well or 1024-well plates) 204 using, for example, bead picking technology described in the just-referenced U.S. Patent application. The use of ‘big bead’ resin (having a diameter of about 500 μ) allows distribution of one bead per well.

The peptoids may then be then dried, cleaved from the resin (e.g., with about 20 – 95% TFA in dichloromethane; 95% being typical), re-dissolved in acetonitrile/water, filtered or centrifuged, dried and re-dissolved in 50% DMSO / 50% Buffer. Other solvent / buffer systems can also be used such as 50% DMF or 50% NMP with the remainder being PBS, TBS, Borate, Tris-HCl, etc. In some cases, a reducing agent such as tris-carboxyethylphosphine (TCEP) may also be added to the wells of the spotting plate to inhibit oxidation of the thiols on the peptoids so that they retain their reactivity toward maleimides on the substrate surface. In cases where TCEP is added to the wells, phosphate buffers such as PBS are to be avoided, and buffers such as TBS would be preferred. This final solution is ready for spotting.

In one embodiment of the present invention, the concentration of peptoids in the wells of the spotting plate should be in the range of about 0.5 – 2 mM, where 2 mM is preferred. For spotting onto the prepared planar solid support 206, the peptoids are filtered or centrifuged, dried, and resuspended, as described in further detail below. It is useful for the material spotted on the slide be in an excess with respect to the amount of probe in solution to be applied to the microarray. This is an amount that gives a saturating fluorescence signal, so that changes in signal intensity are due substantially to protein levels.

sub B2
The anchoring and linker groups may be attached to the peptoid at either the C- or N-terminus. They can be attached either as a submonomer (e.g., as described in U.S. Patent No. 5,877,278 and above-referenced co-pending U.S. Patent Application Serial No. _____ (Attorney Docket No. 16708.001)) during the peptoid synthesis as described above in the patent documents incorporated by reference, or with in situ activated amino acid coupling steps, as a modification of the peptoid after synthesis, according to procedures known to those of skill in the art.

For N-terminal attachment, the peptoid may be synthesized on a resin and then the linker and the anchor groups may be added before the entire molecule is cleaved. For example, peptoids may be prepared on Rink amide polystyrene resin as illustrated and described in co-pending Application No. 09/704,422 previously incorporated by reference herein. The synthesis procedure is also reported in Figliozzi, G. M., Goldsmith, R., Ng, S. C., Banville, S. C., Zuckermann, R. N. *Methods Enzymol.* 1996, 267:437-447, the disclosure of which is incorporated by reference herein for all purposes. Also, before cleavage, one or more (e.g., two to four, preferably four) submonomer hydrophilic linker groups (e.g., methoxyethylamine) may be added to the peptoid N-terminus. Then, a trityl-protected cysteamine is added in order to provide a thiol anchoring group. The peptoid with attached linker and anchoring groups may then be cleaved from the resin, for example using 95 % TFA (v/v) in dichloromethane (CH₂Cl₂). The resulting solution is then ready for application (spotting) onto microarray slides in accordance with the present invention.

In another embodiment, an FMOC-protected beta-alanine is attached to the N-terminus of the peptoid by in situ activation with Hobt and DIC, as is known to those skilled in the arts of peptide synthesis. The beta alanine functions as an adaptor molecule between the peptoid and peptide linker. After attachment of beta-alanine, an FMOC-protected amino acid on a peptide (e.g., glycolic) linker, for example, is attached to the N-terminus of the beta-alanine. Finally, biotin is added to the N-terminus by peptide coupling.

sub B3
The thiol or biotin anchoring group can also be attached to the end of the peptoid at the C-terminus. For example, 4-(diphenylhydroxymethyl)benzoic acid (available from Fluka) is treated with cystamine hydrochloride in the presence of an acid catalyst. Next the resulting amine is protected as the N-(9-fluorenylmethoxycarbonyl) (N-FMOC) derivative, and the resulting FMOC-NH-CH₂CH₂-S-Tr-COOH is coupled to aminomethyl-Big Beads (400-500 microns, Polymer Labs). The peptoid is synthesized on the deprotected amine as described

B3
~~above, and treatment with TFA results in cleavage of the thiol-modified peptoid from the resin, while leaving the trityl protecting group on the resin. Such procedures are described in the above-referenced co-pending U.S. Patent Application Serial No. _____ (Attorney Docket No. PP16708.001)~~

5 C. Contacting the Solid Support with Protein-Binding Agents

Following preparation of the substrate and protein-binding agents, as described above, two or more different protein-binding agents of interest that are to be bound to the surface to produce the array are contacted with the functionalized, or otherwise prepared (cleaned, oxidized, etc.) substrate surface. By contact is meant that
10 the binding agents are brought into proximity with the surface such that they become substantially stably attached or bound to the surface of the substrate layer.

In contacting the binding agents with the substrate surface, any convenient means for contacting the surface with the binding agents which results in the desired pattern of binding agent spots, as described above, may be employed, e.g., by
15 spotting. As mentioned above, the term contact is used herein to refer to any method that brings the binding agent within close proximity of the support surface. Generally, an aqueous solution (e.g. water, water/organic solvent (such as 50/50 water/DMSO, or the like) of the binding agent is employed during contact where the solution may comprise one or more components in addition to water and the binding
20 agent, e.g., buffering agents, salts, and the like. Other systems include 50/50 NMP/TBS, DMF/TBS, NMP/PBS, DMF/PBS, or all these solvents together with water. The 50/50 mix can also be adjusted. For example, when a higher percentage of the aqueous solution is used, the drop sizes can be smaller because of the higher surface tension of the solution. Drop size (and therefore density) may be controlled to
25 some extent in this manner. Typically, contact is achieved by depositing solutions of the different binding agents onto discrete locations of the support surface, such that each different type of binding agent is deposited onto its own unique location on the substrate surface.

The binding agents may be deposited onto the support surface using any
30 convenient means, e.g., by pipetting. A number of devices and protocols have been developed for depositing aqueous solutions onto precise locations of a support surface and may be employed in the present methods. Such devices include "ink-jet" printing devices, mechanical deposition or pipetting devices and the like. See e.g., U.S. Patent Nos. 4,877,745; 5,338,688; 5,474,796; 5,449,754; 5,658,802; 5,700,637;
35 and 5,807,552; the disclosures of which are herein incorporated by reference. Robotic

devices for precisely depositing aqueous volumes onto discrete locations of a support surface, i.e., arrayers, are also commercially available from a number of vendors, including: Genetic Microsystems; Molecular Dynamics; Cartesian Technologies; Beecher Instruments; Genomic Solutions; and BioRobotics. Alternatively, bubble jet technology recently described by Okamoto, Suzuki and Yamamoto, *Nature Biotechnology*, vol. 18 (April, 2000), 438, may be used.

As noted above, an important feature of a process in accordance with the present invention is that the reaction between the anchoring group on the protein-binding agent and the substrate surface must be sufficiently facile so that it is complete within the average lifetime of a droplet that is deposited by the robotic array spotter onto the surface. For example, if the surface is functionalized and displays a maleimide, a suitable anchoring group is a thiol and approximately 15-20 minutes at about 60% humidity are required for completion of the binding reaction. As noted above, other surface display/anchor combinations are possible, including those forming stable, yet non-covalent bonds, such as avidin and biotin.

D. Blocking the Chip

After spotting of the peptoid library onto the array substrate (chip), the remaining, uncoated surface of the chip may be functionalized with a molecule that displays a hydrophilic terminus. These hydrophilic termini are anticipated to reduce or eliminate non-specific binding of proteins in the complex mixture. The hydrophilic portion may consist of alcohols, sulfoxide, carbohydrates, acrylamides, with hydrophilic termini such as alcohols, carbohydrates, amino acids, sulfoxides, acrylamides, and ethers or other low-protein binding group. The hydrophilic display molecule is anchored to the chip in the same manner as the peptoid that has already been spotted. For example, chips spotted with the peptoid library may be chemically blocked with cysteine, mercaptoethanol or other suitable hydrophilic thiol. The chips may also be blocked with protein such as 2% BSA / PBS, 10% non-fat dry milk or 1% casein for at least 1 hour, rinsed with water and dried. Other possible blocking agents are noted above. The blocking agents may be applied to the chips in ways well known to those of skill in the art, such as by dipping the chips in a solution of a blocking agent, by painting the surface of the chips with a blocking agent solution, or by spin-coating.

E. Summary

Sub B4

Figs. 4A and 4B briefly illustrate processes for making protein-binding agent arrays for some embodiments of the invention in accordance with the procedures described above. In Fig. 4A, a process (400) for making an array in which protein-binding agents are bound directly to the inorganic surface of a bare planar substrate is depicted. A planar substrate 412 with a gold or aluminum surface is provided (410). The surface is prepared for binding (cleaned) as described above. Protein-binding agents 422 with a thiol anchoring group 434 are spotted onto the substrate 412 (420). Once binding of the protein-binding agents is complete, a blocking agent 432, namely a hydrophilic group, such as an alcohol, or a protein is applied to the surface of the substrate 412 where no protein-binding agent 422 is bound (430).

In Fig. 4B, a process (450) for making an array in which protein-binding agents are bound to the surface of a planar substrate via an organic surface layer is depicted. A planar substrate 462 with a gold or aluminum surface is provided (460). The surface is prepared for binding by applying a functionalized amino-modified thiol or amino modified silane layer 464, as described above. The layer 464 includes a thiol or silane functionality 466 which binds to the gold or aluminum surface of the substrate 462 and a binding functionality 468, such as maleimide, for a subsequently bound protein-binding agent. Protein-binding agents 472 with anchor group functionality 474 complementary to the substrate surface layer binding functionality 468 are spotted onto the substrate (470). Once binding of the protein-binding agents is complete, a blocking agent 482, namely a hydrophilic group, such as an alcohol, or a protein is applied to the surface of the substrate where no protein-binding agent 472 is bound (480).

3. Methods of Using the Protein-Binding Agent Arrays of the Subject Invention

The subject arrays find use in a variety of different applications in which binding events between the surface bound binding agents of the array and analyte(s) of interest in a test sample are detected. In other words, the arrays of the subject invention find use in binding assays. In such applications, the support bound binding agent generally acts as a "target" for the analyte "probe" in the test sample. The analyte probe is typically labeled, e.g., where the label may be a directly detectable label (e.g., radioactive isotope, fluorescent label, chemiluminescent label, etc.) or an indirectly detectable label (e.g., member of a signal producing system, such as a ligand for a labeled antibody, where the label may be enzymatic which converts a

substrate to a chromogenic product, etc., where the labeled antibody may be a secondary labeled antibody) so that binding events may be readily detected.

In particular, arrays in accordance with the present invention are useful in performing proteomic analyses of complex protein samples. As used herein, proteomics is the separation and/or quantitation and/or identification of one or more proteins in a sample. The sample may be derived from a cell (e.g., the cell's cytosol, membrane or extra-cellular proteins), tissues (e.g., dissected or laser-microdissected), body fluids (such as urine, blood spinal fluid) or any other sample containing proteins. The results of such separation/quantitation/identification may produce novel protein targets for drug screening, proteins for diagnostics, or novel synthetic ligands for assays or protein purification. The arrays may very effectively be used in differential protein binding assays. For example, two (or more)-color fluorescent labeling of complex protein mixtures, and the analysis of differential protein binding to the array by fluorescence imaging may be conducted. As described below, the arrays may be used in conjunction with other techniques to identify, sequence and structurally characterize differentially expressed proteins or peptides of interest. The arrays may be run in parallel with DNA arrays and the differential binding results compared to identify correlations between gene activity and protein expression. Also, mixed arrays, wherein the molecules making up an array includes antibodies, etc. may be prepared and used to conduct binding assays.

A variety of techniques can be used to conduct differential binding assays using arrays in accordance with the present invention ("proteomic microarrays"). Some of these techniques, as used in embodiments of the present invention, are described below:

A. Protein Labeling

Complex protein samples are labeled using standard techniques, many of which have been developed for 2-D gel analysis of protein mixtures. For example, sample A may be labeled with an amine reactive Cyanine 3 dye ("Cy 3") ($\lambda_{\text{ex}} = 550\text{nm} / \lambda_{\text{em}} = 570\text{nm}$), and sample B is labeled with an amine reactive Cyanine 5 dye (Cy 5) ($\lambda_{\text{ex}} = 650 / \lambda_{\text{em}} = 670 \text{ nm}$) (dye reagents available from Amersham-Pharmacia). Samples A and B may be, for example, from normal or diseased, treated or untreated, etc., tissues or cell lines, respectively. The unreacted dye may be separated from the labeled protein using standard methods such as gel filtration, dialysis, etc. Of course, as noted above, a variety of different labels, as are well known to those of skill in the art, including, but not limited to,

tetramethylrhodamine—isothiocyanate (TRITC), fluorescein-isothiocyanate (FITC) , and succidimidyl ester derivatives, thereof, or any other dye molecule that may be reacted to proteins via amino acid side chains such as amine side chains (lysine), thiol side chains (cysteine) or other suitable functional group.

5 B. Binding Assay and Chip Readout

Labeled protein samples are incubated with the proteomic microarray chip for periods of time, and under a variety of conditions of pH, salt content and temperature anticipated to modulate the affinity of various proteins to the elements of the array. Generally, the samples are contacted with the microarray by introduction of an appropriate volume of the fluid sample onto the array surface, where introduction can be flooding the surface with the sample, deposition of the sample onto the surface, e.g., with a pipette, immersion of the entire array in the sample, and the like. In many embodiments, the solution is deposited onto the surface and then sandwiched beneath a cover slip or in a sealed chamber.

For example, a 25 μ L – 100 μ L (typically 50 μ L) aliquot of each probe solution may be applied to the surface of a typical microscope slide-sized chip, and a cleaned coverslip placed on top, forming a sandwich of the probe solution on the chip surface. The protein solutions may then be co-incubated with the chip for at least 1 hour, or overnight. After incubation, the coverslip is removed and the chip is washed, for example, in 1X PBS / 0.05% Tween or other suitable buffer containing surfactant. The chip may be washed using a variety of conditions that decrease or increase stringency. These conditions can again be customized to allow, for example, retention of only the most strongly bound proteins. Or, as the case may warrant, less stringent washing may be used to allow visualization of comparatively weaker bound proteins. The choice is likely to be determined by the complexity and diversity of the peptidomimetic (e.g., peptoid) array that is displayed on the chip and the nature of the protein mixture. The washed chips are then dried, for example, under a stream of Argon or Nitrogen.

After suitable washing, the chip is read in an array scanner, such as are well known in the art. The ratio of Cy 3 to Cy 5 for each spot is determined using commercially available software. Spots that show a ratio considerably greater than or less than one are observed, and deemed to be “differential”.

Fig. 5 briefly illustrates a process for conducting a differential proteomic binding assay using protein-binding agent arrays for one embodiment of the invention

in accordance with the procedures described above. In Fig. 5, the process (500) begins with the procurement of two biological samples to compare, e.g., an “untreated” cell line 502a and a “treated” cell line 502b. Cell lysates 504a,b are isolated from the cell line samples. The lysates are labeled, for example, the “untreated” cell lysate 504a is labeled with a fluorescent green dye while the “treated” cell lysate 504b is labeled with a fluorescent red dye. The labeled samples 506a,b are then co-incubated on a protein-binding array chip 508 in accordance with the present invention, e.g., an array of peptoids. The protein in the samples can either be denatured or native. For example with the addition of 1-2% SDS the proteins in the samples may be denatured and clusters or hydrophobic interactions minimized or eliminated. Alternatively, the clusters, which may be important in elucidating protein-protein binding pathways, and proteins may be kept in their native states and the results studied. The chip is then read in an array scanner.

C. Library Sequencing

The sequence of the peptidomimetic that binds a differentially expressed protein may be determined by library sequencing techniques. This can be achieved, for example, by MS/MS methods that allow fragmentation of the peptoid along the amide backbone. These methods are described in co-pending U.S. Patent Application Serial No. 09/580,380, the disclosure of which is incorporated herein by reference for all purposes (*see also* PCT Publication No. WO 00/72004). From the mass of the fragmentation product, the structure and sequence of the isolated peptoid can be determined.

D. Post-Array Processing: Protein Isolation, Purification and Identification

Once a protein or set of proteins is determined to be differential between samples A and B, it can be isolated by preparing chromatographic supports composed of the same peptidomimetic (e.g., peptoid) identified on the chip. Peptoid-based chromatographic supports, their preparation and their use are described in U.S. Patent Application Serial No. 09/704,422, previously incorporated by reference herein.

In accordance with one embodiment of that disclosure, a peptoid sequence that produces a differential on the chip may be synthesized on hydrophilic resins that are hydrophobically “masked” during peptoid synthesis. After synthesis, the resin is “unmasked” to reveal hydrophilic groups compatible with biological solutions. Such a resin is immediately available for protein binding / isolation experiments.

Once the protein is isolated, it's sequence can be determined using standard techniques such as MALDI / TOF mass spectrometry or trypsin digests.

Fig. 6 briefly illustrates aspects of post-array processing in accordance with the procedures described above and below. In Fig. 6, the process (600) begins with the preparation of chromatographic separation columns 602 using peptidomimetic agents 601, e.g., peptoids, identified as of interest in a proteomic differential binding assay conducted using a proteomic microarray in accordance with the present invention. An aliquot of the complex sample originally run on the microarray is then run through the column. The protein of interest preferentially binds to the column and is thereby separated from other components of the sample. The bound protein is eluted and may then be used in further analyses 604, such as protein sequencing, tertiary structure determination, etc. In addition, data relating to the identification of the protein may be entered into bioinformatics databases for further research.

Alternatively, the same peptoid that bound the differentially expressed protein could be spotted repetitively on a chip and incubated with the an aliquot of the same lysate. The same protein should bind to the array, but in a much larger area than just the one spot on the original chip. Laser desorption mass spectrometry can then be used to sequence the protein directly from the chip.

E. Other Applications

The anticipated uses of proteomic microarray chips in accordance with the present invention are broad. In general, the applications have in common the identification of a protein or set of proteins that are over-expressed or under-expressed in one complex mixture relative to another (or present/absent, such as in the case of a diagnostic protein). Those skilled in the art will recognize that embodiments of the present invention are compatible with a wide variety of assay formats including sandwich assay, such as ELISA.

As described above, proteomic microarrays may be used to determine differential expression of proteins in complex solutions by alternatively labeling (e.g., Cy 3 for one sample and Cy 5 for another) the two or more protein solutions to be compared. The chips may be used to find novel protein targets for later high throughput screening assays. In another particularly powerful application, the methodology may be used to purify a recombinant protein that is overexpressed in a particular host such as yeast or baculovirus. The sample that contains the expressed protein is compared to the sample that does not by co-binding the alternatively labeled

samples on the chip, and looking for differentials. The procedure identifies peptoids on the array that bind with reasonable affinity and specificity to the expressed protein. These same peptoids are then used for generating chromatographic resins for the isolation and purification of the recombinant protein of interest.

5 In an analogous manner, the proteomic microarray chips may be used to find protein markers in plasma or serum that may be diagnostic of particular disease states such as cancer, HIV, or diabetes, or to find novel targets for drug screening. Also, once a set of ligands has been identified for particular groups of proteins, it is possible to monitor the expression levels of these proteins to decipher mechanisms of drug
10 action. In that regard, the new ligands may be used as probes of protein abundance, analogous to the ways in which antibodies are currently used to determine protein abundance. In addition, by examining the proteins from virulent and non-virulent strains of bacteria or viruses, one can determine unique virulence factors that result in infectious disease. Once these virulence factors are identified, these proteins can be
15 used as targets for screening new anti-bacterial or anti-viral drugs. Alternatively, the chips provided by the present invention may also be used to discover a variety of peptide, antigen, or protein mimetics. For example, novel mimetic cellular adhesion molecules, mimetic drug candidates, or protein mimetics that may be used as chromatographic supports. In addition, the material spotted on the chip could itself be
20 potential drug candidates or function as an initial scaffold for designing new drug candidates. In such embodiments, high throughput assays to identify potential therapeutic agents can be done directly on the chip.

In one embodiment, the proteomic microarrays of the invention are run in parallel with DNA arrays, and the differential binding results derived from each are
25 compared to identify correlations in gene activity and protein expression. For example, differential binding assays are conducted for complex biological samples on both proteomic and DNA arrays. Separate aliquots from the samples are labeled and contacted with a proteomic microarray in accordance with the present invention and a DNA microarray, such as are well known in the art. The differential protein
30 expression evidenced by the binding results on the proteomic array when compared with those for the DNA array may elucidate relationships between protein expression and gene families whose activation is required for that expression.

In another embodiment of the present invention, a "mixed array" chip of the present invention is provided. Peptoid or other peptidomimetic binding elements
35 attached to the chip surface may be mixed with antibody or protein binding elements

such that both types of binding elements are present on the same chip. The antibodies can serve as positive controls, or as a means of monitoring the levels of specific proteins in the mixture being analyzed. The peptoids would provide data on differentials in unknown proteins, whereas antibody spots on the chip would provide data on differential levels of known proteins. For example, if a cell is treated with a certain drug, the protein levels might change up or down. If those proteins have known antibodies, it is possible to monitor the change of these with antibody differentials, while at the same time, look for changes in unknown proteins, with the peptoid differentials.

In one example of this technique, antibody solutions are prepared in the same microtiter plates as the peptoid solutions, but in different wells. The peptoids are already functionalized with thiol anchoring groups. The antibodies are reacted with a reducing agent to reduce the disulfide bonds in the antibody hinge and Fc region (for example, as described in Levison, M.E., et al, (1969), *Experientia*, vol 25, 126-127 and Blauenstein, P., et al, (1995), *Eur. J. Nuclear Med.*, vol 22, 690-698), thus producing a thiol on the antibody. This allows for spotting of both peptoid and antibody in the same spotting session, thereby creating a "mixed" chip. Alternatively, the antibodies may be bound to immobilized Protein A or Protein G on chip surfaces, while peptoids are attached to avidin or streptavidin, presenting a mixed surface for display. Or, antibodies may simply be biotinylated and spotted together with biotinylated peptoids or peptides onto avidin-coated chips.

Another technique that may be combined with the proteomic microarray techniques of the present invention is the MS/MS macromolecular structural analysis technique described in above-referenced U.S. Patent Application Serial No.09/580,380, previously incorporated by reference herein. In this way, the combination of techniques can be used to identify a protein of interest, enrich and isolate it, sequence the protein, and elucidate aspects of its tertiary protein structure.

Data relating to the identification and post-array processing of proteins of interest may also be entered into bioinformatics databases. The data may be correlated with other biological data therein for further research.

4. Kits

Also provided by the subject invention are kits for performing proteomic binding assays using the subject arrays. Such kits according to the present invention will at least include an array according to the invention. The kits may be configured

for analytical or diagnostic purposes, and may further include one or more additional reagents employed in the method for which the array is intended. For example, the kit may include various receptacles, labels, buffer solutions, tools and any other material necessary to conduct a proteomic binding assay. Kits in accordance with the present invention may also be configured to receive samples for analysis and thereafter perform the steps necessary for a binding assay in accordance with the invention without further user manipulation.

EXAMPLES

The following examples provide details concerning the synthesis and characteristics of the proteomic arrays in accordance with the present invention, their components, and applications. It should be understood the following is representative only, and that the invention is not limited by the detail set forth in these examples.

Example 1: Preparation of microtiter plates containing one compound per well

Preparation of one (1) 96-well plate containing 96 single compounds (0.8 mM) in a 1:1 DMSO/PBS solution: A siliconized vial (8 ml, 2 dram) was loaded with a one compound/bead library synthesized on Rink polystyrene macro-bead solid support (66 mg, ~1320 beads, 40 nanomole/bead, 0.75 mmol/g, 425-500 um, Polymer Laboratories) that contains a total of 121 possible compounds. The beads were swollen in dichloroethane (DCE, 4 ml) for 24 hours and sieved over stainless steel mesh (600 um). Using the resulting dichloroethane bead slurry, 96 beads were individually picked and relocated into a polypropylene 96-well plate (200 µL, conical bottom) in order to obtain one bead per well over 96 wells. The resulting 96-well plate, (the 'grandmother' plate), was transferred to a speed-vac evaporator Savant AES200 equipped with a 96-well plate carrier rotor and the remaining DCE was removed from the plate. A cleavage cocktail (TFA/TES/H₂O/DCE, 46:2:2:50, 75 µL) was then added to each bead containing well in the plate. After an hour the plate was transferred to the speed-vac evaporator to remove most of the volatiles from the plate. Each well was treated with acetonitrile (CH₃CN, 10 µL) and agitated for 10 min using a microtiter plate shaker. The plate was transferred to a speed-vac evaporator and the remaining volatiles were removed from the plate for 10 minutes. Every single compound in each well was dissolved in dimethylsulfoxide (DMSO, 25 µL) and relocated in a 96-well filter plate made of polystyrene, equipped with a polypropylene membrane (0.45 µm).. The 96-well filter plate was stacked over a polypropylene 96-well plate (200 µL, conical bottom) and the assembly was

transferred to a Beckman GS-6R centrifuge programmed for 5 minutes at 3000 RPM (~2000G) at 15 °.

This procedure provided the 'mother' plate composed of filtered 96 DMSO solutions (1.6 mM) of one compound in 96 wells. To form the 'daughter plate' (used for spotting), an aliquot of every DMSO solution (5 µL) was transferred to a polypropylene 96-well plate (200 µL, conical bottom). Then, every well in the daughter plate was mixed with degassed solution PBS (5 µL) and stored at -20°C. Alternatively, each compound in each well may be dissolved in 20 µL of 50% DMSO / water. In the case of a plate with a conical bottom, the plate may be centrifuged at 1,800 rpm for 5 minutes to immobilize the bead at the bottom of the conical well, followed by relocation of 10 uL of the clear supernatant into a daughter plate that is ready for spotting.

The same can be done for 384 beads in 384 well plates, or 1,024 beads in 1,024-well plates, etc. For spotting, any suitable spotting device can be used, such as a Molecular Dynamics Generation II (for 96-well) or Generation III (384-well) spotters.

Example 2: Functionalization of solid supports for spotting

Gold or aluminum reflective microscope slides were used as substrates for spotting a library of peptoids. In one example, the peptoid is functionalized with a thiol endgroup and the surface is functionalized with a maleimide. Upon spotting, the thiol on the peptoid reacts with the maleimide on the surface to form a covalent thioether attachment. Gold surfaces activated with maleimide were prepared as follows: 1) Gold-coated microscope slides (1200 Angstroms Au, 30-50 Angstroms Ti or Cr) were cleaned with Chromic acid cleaning solution for 15 minutes and rinsed with HPLC grade water. 2) Gold slides were dipped into 1-5 mM amino-modified thiol (1-Mercaptoundecyldiethoxyamine; a C-11 alkyl, two ethylene oxides, and an amine; alternatively, C-2 to C-20 alkyl groups and/or ethoxy or triethoxy groups could be used in such a compound) for 1-24 hours at room temperature or at 45 or 60°C. The slides were rinsed four times in absolute ethanol and dried under a steam of Nitrogen. 3) The amino-modified gold slides were dipped into a solution (50-100 uM) of succinimidyl 4-(N-maleimidomethyl)-cyclohexane-1-carboxylate (SMCC) to render a surface that presents the maleimide functional groups.

Aluminum slides were made or purchased from Amersham-Pharmacia, coated with a 1000 – 1400 Angstrom layer of silicon oxide, and a layer of aminopropysilane

(APS). The amino-modified Al surfaces are functionalized with SMCC as described above.

In some cases, after spotting the peptoid library, the aluminum slides were etched in a solution of 0.1% SDS / 5X SSC at 60 degrees C for 2 hours. The etched slides exhibit a reduced oxide layer thickness (200 – 900 Angstroms) to allow amplification of both the Cy5 and Cy3 signals.

Example 3: Derivatization of Aluminum / SiO₂ surfaces with avidin

Aluminum slides coated with aminosilane were dipped into a solution of NHS-LC-LC-biotin (“LC” refers to 6-aminohexanoyl and “NHS” refers to N-hydroxysuccinimidyl) (commercially available from Pierce), depicted in Fig. 7A, that was 0.39mM in PBS buffer. The slides were coated for 1.5 hours with shaking at 80 rpm. After attachment of biotin, the slides were rinsed with water, then dipped in a solution of 1ug/ml - 1mg/ml avidin, streptavidin or neutravidin in PBS buffer for 2 hours, stirring at 70rpm. The slides were rinsed with water and ready for spotting biotinylated peptides or peptoids. Fig. 7B depicts a representation of an avidin-derivatized aluminum slide spotted with a biotinylated protein-binding agent in accordance with one embodiment of the present invention.

The various surface modification steps was followed using ellipsometry to note the thickness changes. A thickness change increase of 40-45 angstroms was reproducibly recorded after the addition of avidin to the surface layers.

Example 4: Protein labeling

Protein solutions were adjusted to a concentration of 1 mg/mL in 0.1 M sodium carbonate, pH 9.3 and a volume of 0.1 – 1 mL, and mixed with bifunctional or mono-functional amine-reactive cyanine dye (Cy3 or Cy5, Amersham Pharmacia). The protein was purified from the unreacted dye by size exclusion chromatography using a Sephadex G-25 packing in a 5 cm long, 1.7 cm diameter column with a 1 mL load, 0.5 mL fractions, and a dilution factor of 3.5.

Fig. 8 provides a graph of the results illustrating that size exclusion chromatography of the components efficiently separates the labeled protein from the unreacted dye. The graph shows the different elution profiles for the protein (BSA standard) and dye molecules.

Example 5: Chip binding experiments

In some instances, the chips spotted with the peptoid library are chemically blocked with cysteine, mercaptoethanol or other suitable hydrophilic thiol. The chips are blocked with protein such as 2% BSA / PBS, 10% non-fat dry milk or 1% casein for at least 1 hour, rinsed with water and dried. The labeled protein probe solution is diluted accordingly (typically to 20 – 1000 ng total protein) with the blocking solution (for the etched Al slides, detection limits of a few picograms have been observed). A 30 – 100 μ L aliquot of the probe solution is applied to the chip surface, and a clean coverslip placed on top, forming a sandwich of the probe solution on the chip surface. The protein solution is incubated with the chip for at least 1 hour. The coverslip is removed in 1X PBS / 0.05% Tween or other suitable buffer containing surfactant. The chip is then washed in 1X PBS / 0.05% Tween or other suitable buffer / surfactant system. The chips are further rinsed with water, dried under a stream of Argon or Nitrogen and scanned.

Example 6: Sample Chip

A microarray chip bearing a library of 1,000 peptoid-based protein-binding agents was prepared in accordance with the present invention. Streptavidin was spiked into cells induced to express proteins in the wnt pathway and into control cell lysate and the entire mixture labeled with Cy3 (control) or Cy5 (wnt). The mixture was then incubated with the chip (6.3 ng total protein/chip; 31 pg streptavidin/chip), which had a biotin-displaying peptoid in it's upper left corner. Figs. 9A and 9B show 1/6 of the incubated chip. As shown, in addition to the biotin peptoid spots, many other peptoid spots are binding to the protein in the lysate. By comparing the relative signals from the Cy5 and Cy3 channels, differentials can be identified. Fig. 9C shows 1/2 of a streptavidin-only control chip (31 pg streptavidin/chip). Biotin peptoid spots are visible in the upper left corner.

Example 7: Demonstration Chip

As illustrated in Fig. 10, hexameric peptides of different and known affinity to the "anti-glu" antibody were synthesized with a biotin anchoring group and heptameric glycyl linker. To prove the concept of the present invention, the biotinylated peptides were spotted onto avidin-treated slides as described in the present disclosure. The slides were blocked with casein and probed with Cy5-labelled antiglu antibodies. The gradations in signal intensity correlate with the known

b5
differences (e.g., measured by direct ELISA) in affinity between the peptides and their cognate antibody probe.

Example 8: Comparison of Surface Attachment Systems

5 Fig. 11 depicts the dependence of signal strength on the mode of surface attachment of a protein-binding agent to a substrate. In this example, signal increases of 1000X are observed for biotin / avidin immobilization compared to thiol / maleimide (signal results depicted above surface attachment description and molecular structure).

Example 9: Preparation of Protein-Coated Slide Surface for Antibody Display

10 To prepare Protein A- or Protein G-modified slide surfaces, a slide coated with avidin (prepared as described above) was immersed in a solution of biotinylated Protein A or Protein G (0.5 - 1mg/mL in PBS buffer, purchased from Pierce, product numbers 29989zz and 29988zz) for 2 hours at room temperature. The slides were then rinsed with de-ionized distilled water and blown dry with Nitrogen or Argon.

15 Conclusion

20 Although the foregoing invention has been described in some detail for purposes of clarity of understanding, it will be apparent that certain changes and modifications may be practiced within the scope of the invention. For example, while the present disclosure emphasizes the use of peptoids as the peptidomimetic segment of the protein-binding array elements, it should be understood that the scope of the invention is not so limited and other molecules having the appropriate properties and function as described herein may also be used. It should also be noted that there are many alternative ways of implementing both the process and apparatus of the present invention. Accordingly, the present embodiments are to be considered as illustrative and not restrictive, and the invention is not to be limited to the details given herein.

What is claimed is: